

REMARKS ON THE BRST-COHOMOLOGY
FOR $c_M > 1$ MATTER COUPLED
TO “LIOUVILLE GRAVITY”ADEL BILAL[★]*Joseph Henry Laboratories
Princeton University
Princeton, NJ 08544, USA*

ABSTRACT

We describe the (chiral) BRST-cohomology of matter with central charge $1 < c_M < 25$ coupled to a “Liouville” theory, realized as a free field with a background charge Q_L such that $c_M + c_L = 26$. We consider two cases:

- a) matter is realized by one free field with an imaginary background charge,
- b) matter is realized by D free fields: $c_M = D$.

In case a) the cohomology states can be labelled by integers r, s of a rotated $c_M = 1$ theory, but hermiticity imposes $r = s$. Thus there is still a discrete set of momenta $p_M(r, r)$, $p_L(r, r)$ such that there are non-trivial (relative) cohomology states at level r^2 with ghost-numbers 0 or 1 (for $r > 0$) and ghost-numbers 0 or -1 (for $r < 0$). The (chiral) ground ring is isomorphic to a subring of the $c_M = 1$ theory which is $(xy)^n$, $n = 0, 1, 2, \dots$, and there are *no* non-trivial currents acting on the ground ring.

In case b) there is no non-trivial relative cohomology for non-zero ghost numbers and, for zero ghost number, the cohomology groups are isomorphic to a $(D - 1)$ -dimensional on-shell “transverse” Fock space. The only exceptions are at level 1 for vanishing matter momentum and $p_L = Q_L(1 + r)$ with $r = \pm 1$, where one has one more ghost-number zero and a ghost-number r cohomology state.

All these results follow quite easily from the existing literature.

★ on leave of absence from Laboratoire de Physique Théorique de l’Ecole Normale Supérieure,
24 rue Lhomond, 75231 Paris Cedex 05, France (unité propre du CNRS)

1. Introduction

Whereas the BRST-cohomology for critical strings has been computed already some time ago [1-4], it is only quite recently that it has been worked out for non-critical strings with matter central charge $c_M \leq 1$ [5,6]. The basic assumption is that 2D gravity coupled to $c_M \leq 1$ matter can be described in the conformal gauge by coupling the matter theory to a free Liouville field with imaginary background charge. A very interesting structure has emerged where cohomology states with non-zero ghost-numbers lead to a ground ring and those with zero ghost-numbers to symmetries of that ground ring [7]. This structure has been used to obtain non-trivial information about correlations functions [8].

This analysis was restricted to $c_M \leq 1$. It is very interesting to know what happens if $1 < c_M < 25$. In fact, the results follow quite easily from the analysis of refs. 4, 5 and 7, but since we are not aware in the literature of any explicit discussion of this BRST-cohomology for $1 < c_M < 25$, we found it nevertheless worthwhile to spell out the results in this note. We will consider two cases:

a) The matter sector is realized by one free field with an imaginary background charge Q_M : $c_M = 1 - 12Q_M^2 > 1$.

b) The matter sector is realized by D free fields (with euclidean or minkowskian signature): $c_M = D$.

Of course we could also consider the more general case of D free fields, one of which has a background charge. However, by a rotation in field space we may reduce this again to case b), and we will comment briefly on this situation after analysing case b).

The results we find are those announced in the abstract. Note that we always refer to the relative cohomology H_{rel}^n (for one chiral sector[†]), i.e. we only consider states annihilated by the anti-ghost zero-mode b_0 . The absolute cohomology then is simply $H_{\text{abs}}^n \simeq H_{\text{rel}}^n \oplus c_0 H_{\text{rel}}^n$.

In a certain sense, these results are quite disappointing, since one might have hoped that some particular BRST-cohomology might occur in the special dimensions put forward in refs. 10. Although the approach of refs. 10 advocates the use of two sets of oscillators in the

[†] The non-chiral, i.e. closed string cohomology involves some subtleties [9], and we will not discuss it here explicitly.

same Fock space of the Liouville sector, related by a complicated transformation, (instead of using screening operators), the Fock spaces used (for each p_L) are the same as in our analysis. Thus our results are applicable and we conclude that, as far as only the BRST-cohomology is concerned, nothing special happens at the special dimensions of refs. 10.

In the present note, we will try as much as possible to remain within the Fock space approach of ref. 5 which we believe to be the most physical one (although some arguments could be short-cut by using the results of ref. 4).

2. Matter is one free field with an imaginary background charge Q_M (case a).

Formally, the analysis of the BRST-cohomology proceeds exactly as in ref. 5, henceforth referred to as BMP,[‡] for the product of the matter and Liouville Fock spaces $\mathcal{F}(p_M) \otimes \mathcal{F}(p_L)$, but now the matter background charge is imaginary, whereas in BMP it was real. However, as far as the BRST-cohomology is concerned, the whole analysis is insensitive to the actual value of the background charges Q_M, Q_L (only constrained by $Q_M^2 + Q_L^2 = -2$ to ensure $c_M + c_L = 26$). As a result, we know that non-trivial cohomology states appear at level rs (≥ 0) with r, s non-zero integers, iff $p_L = p_L(r, s)$, $p_M = p_M(r, s)$ with

$$\begin{aligned} p_L(r, s) - Q_L &= Q_L \frac{r+s}{2} - iQ_M \frac{r-s}{2} \\ p_M(r, s) - Q_M &= Q_M \frac{r+s}{2} + iQ_L \frac{r-s}{2} . \end{aligned} \tag{2.1}$$

Then there is always one (relative) cohomology state with zero ghost-number and one with ghost-number $\text{sign}(r)=\text{sign}(s)$. (For r or s zero, there is only a ghost-number zero cohomology state: the corresponding Fock space vacuum.) For $c_M \leq 1$, Q_M was real and Q_L imaginary, hence $p_M(r, s)$ was real and $p_L(r, s)$ imaginary, which was appropriate for hermiticity. Now, for $1 < c_M < 25$, we have Q_M and Q_L imaginary and for general r, s we get complex $p_M(r, s)$ and $p_L(r, s)$. This implies that, in general for $p_L = p_L(r, s)$, $p_M = p_M(r, s)$, the matter and Liouville stress tensors are not hermitian separately, only their sum is. We believe, however, that it is a reasonable physical requirement to impose hermiticity separately on the matter

[‡] We use the same notations and conventions as BMP.

stress tensor T_M and on the Liouville one T_L . Let us briefly discuss the consequences of this requirement.

Consider one free field with background charge Q and Virasoro generators

$$L_n = \frac{1}{2} \sum_{m \neq 0, n} : a_m a_{n-m} : + (p - (n+1)Q) a_n \quad (2.2)$$

where $p \equiv a_0$. Since the dual of the Fock space $\mathcal{F}(p)$ is isomorphic to $\mathcal{F}(2Q - p)$, hermiticity means

$$L_n(p)^\dagger = L_{-n}(2Q - p) . \quad (2.3)$$

Comparing the quadratic terms simply implies $a_n^\dagger = \pm a_{-n}$ while comparing the linear terms implies the following:

$$\begin{aligned} Q \text{ real} &\Rightarrow a_n^\dagger = -a_{-n} , \quad p^* = p \\ Q \text{ imaginary} &\Rightarrow a_n^\dagger = +a_{-n} , \quad p^* = -p . \end{aligned} \quad (2.4)$$

We see that hermiticity requires that p is real if Q is real and p is imaginary if Q is imaginary.

Going back to our case of interest, Q_M and Q_L imaginary, we conclude that p_M and p_L should be imaginary. From eq. (2.1) we see that $p_M(r, s)$ and $p_L(r, s)$ are imaginary only for $r = s$:

$$\text{hermiticity} \Rightarrow r = s . \quad (2.5)$$

Thus, in the hermitian case, we have non-trivial cohomology iff

$$\begin{aligned} p_L &= p_L(r, r) = Q_L(r+1) \\ p_M &= p_M(r, r) = Q_M(r+1) . \end{aligned} \quad (2.6)$$

In these cases the cohomology states (with ghost-numbers 0 and 1 for $r > 0$ and 0 and -1 for $r < 0$, and 0 for $r = 0$) are formally (i.e. when expressed in terms of Q_M, Q_L) the same as those constructed by BMP.

Just as in refs. 7, 11, we may now construct fields $\Psi_{r,r}$ ($r < 0$) forming a (chiral) ground ring[§] and currents $J_{r,r}$ acting on the ground ring. One finds that the (chiral) ground ring is simply isomorphic to the ring of non-negative integers (with the addition)

$$\Psi_{r,r}\Psi_{r',r'} = \Psi_{r+r'+1,r+r'+1} \quad (2.7)$$

(modulo BRST commutators), and that the currents act trivially:

$$J_{r,r}(\Psi_{r',r'}) = 0. \quad (2.8)$$

Probably the easiest way to see that this must be true is to remember[¶] that we may perform a rotation (with real coefficients) in the space of the two fields ϕ_M and ϕ_L to obtain new fields $\tilde{\phi}_M$ and $\tilde{\phi}_L$ such that $\tilde{\phi}_M$ is a $c = 1$ (matter) field and $\tilde{\phi}_L$ a $c = 25$ (Liouville) field. Then eq. (2.1) implies $\tilde{p}_M = \frac{1}{\sqrt{2}}(s - r)$. From our discussion above we know that hermiticity of the $c_M > 1$ matter stress tensor requires $r = s$, which we now interpret as $\tilde{p}_M = 0$. In other words, the cohomology of hermitian matter with $1 < c_M < 25$ is isomorphic to the one of the $\tilde{p}_M = 0$ sector of the $c_M = 1$ theory. Of course, we can rotate back the results of Witten on the ground ring and symmetry currents [7] after restriction to this $\tilde{p}_M = 0$ ($r = s$) sector. The only elements of the ground ring $x^n y^m$ (in the notation of ref. 7) that survive the projection are those with $n = m$, and the projected ground ring is indeed isomorphic to the ring of non-negative integers. The chiral symmetry currents of the $c_M = 1$ theory act as $\frac{\partial h(x,y)}{\partial x} \frac{\partial}{\partial y} - \frac{\partial h(x,y)}{\partial y} \frac{\partial}{\partial x}$, but this can never map $(xy)^n$ to $(xy)^m$. Hence the symmetry currents that survive the projection must all be represented by zero when acting on the projected ground ring. Of course, we may check this directly in the original $c_M > 1$ theory: $J_{r,r}$ and $\Psi_{r,r}$ carry momenta $p_L = Q_L(r+1)$, $p_M = Q_M(r+1)$. Thus e.g., a priori, $J_{0,0}(\Psi_{-2,-2})$ must be proportional to the identity $\Psi_{-1,-1}$. Computing the OPE explicitly we see indeed that the various contributions (proportional to the identity) cancel among each other and we obtain zero.

To conclude, hermiticity restricts us to $r = s$. The chiral ground ring is isomorphic to $(xy)^n$, $n = -(r+1) \geq 0$, while all chiral symmetry currents $J_{r',r'}$ act trivially ($= 0$) on the ground ring.

[§] We only consider the chiral ground ring here. Extending the discussion to the full non-chiral ground ring along the lines of refs. 7 and 9 is straightforward.

[¶] We thank E. Witten for reminding us of this fact.

3. Matter is D free fields: $c_M = D$ (case b).

Consider now D free fields (without background charge) with euclidean or minkowskian signature. As we will see, the signature is without influence on the result of the analysis. Although the results can be quite easily be foreseen from the vanishing theorem of ref. 4, we wish to stay within the framework of Fock spaces as far as possible.

We will label the (real) matter momentum by p with components p_i , $i = 1, \dots, D$ and assume first that $p \neq 0$. Our strategy is to show that we can always chose coordinates such that the BMP analysis becomes particularly simple. In particular we may assume that $p^+ = \frac{1}{\sqrt{2}}(p_1 + ip_2)$ is non vanishing. We further define $a_n^\pm = \frac{1}{\sqrt{2}}(a_n^1 \pm ia_n^2)$.

Recall that non-trivial cohomology is only possible on-shell, i.e. for zero eigenvalue of $L_0 = \{b_0, d\}$, b_0 being the anti-ghost zero-mode and d the BRST operator^{*}. The $L_0 = 0$ condition reads

$$\frac{1}{2}(p_L - Q_L)^2 - \frac{1}{2}Q_L^2 + \frac{1}{2}p^2 + R - 1 = 0 \quad (3.1)$$

where R is the total level operator (matter plus Liouville plus ghosts) and

$$Q_L^2 = -\frac{25 - D}{12}. \quad (3.2)$$

We will consider the relative cohomology which is the cohomology on the subspace of the Fock space annihilated by b_0 (and L_0). On this subspace, the BRST operator $d = c_0 L_0 - b_0 M + \hat{d}$ reduces to $\hat{d}$ (note $\hat{d}^2 = 0$). Following BMP, we break up $\hat{d}$ into three pieces, $\hat{d} = \hat{d}_0 + \hat{d}_1 + \hat{d}_2$, of definite degrees 0, 1 and 2 where we define the degree by

$$\begin{aligned} \deg(a_n^+) &= \deg(c_n) = 1 \\ \deg(a_n^-) &= \deg(b_n) = -1 \quad (n \neq 0) \end{aligned} \quad (3.3)$$

and assign degree zero to all other operators and to the Fock space vacua. The benefit of all this is that

$$\hat{d}_0 = p^+ \sum_{n \neq 0} c_{-n} a_n^- \quad (3.4)$$

is particularly simple, and that in the present case it turns out to be enough to investigate the

* Indeed, if $L_0 \psi = h \psi$ and $d \psi = 0$ then for $h \neq 0$ we have $\psi = (1/h) L_0 \psi = d(b_0 \psi / h)$.

cohomology of $\hat{d}_0$. Indeed, one has the following theorem (BMP) [5] (C_k denotes the subspace of degree k):

Theorem: If for each ghost-number n the cohomology classes $H^n(C_k, \hat{d}_0)$ are non-zero for at most one degree k , then $H^n(C_k, \hat{d}_0)$ and $H^n(C_k, \hat{d})$ are isomorphic.

Now, by the usual trick, the cohomology of $\hat{d}_0$ is very easily obtained by observing that the operator $K = \frac{1}{p^+} \sum_{n \neq 0} b_n a_{-n}^+$ satisfies the following anticommutation relation with $\hat{d}_0$:

$$\{\hat{d}_0, K\} = \sum_{n \neq 0} (a_{-n}^+ a_n^- + n c_{-n} b_n) \equiv \hat{R} . \quad (3.5)$$

By precisely the same argument as for L_0 we now conclude that non-trivial $\hat{d}_0$ -cohomology can only occur for zero eigenvalue of $\hat{R}$. Thus the cohomology states are all vectors in the Fock space that contain neither ghost nor $a^\pm$ excitations. These are ghost number zero states made up from an arbitrary number of Liouville and $a^i, i = 3, \dots, D$ oscillators such that the on-shell condition (3.1) is satisfied. Since all these states have zero degree, the above-cited theorem does apply and the $\hat{d}_0$ and $\hat{d}$ cohomologies are isomorphic. For R a non-negative integer, let $\mathcal{P}_{D-1}(R)$ denote the number of states at level R in a Fock space obtained by applying $D-1$ sets of oscillators a_{-n} to the vacuum. (This is the coefficient of q^R in $\prod_{n=1}^{\infty} (1 - q^n)^{1-D}$.) Let further $\mathcal{P}_{D-1}(R)$ equal zero whenever R is not a non-negative integer. Then we may state the result for the $\hat{d}$ cohomology - which is in fact the relative d cohomology - as

$$\dim H_{\text{rel}}^n(\mathcal{F}(p_L) \otimes \mathcal{F}(p), d) = \delta_{n0} \mathcal{P}_{D-1}(R) \quad (p \neq 0) \quad (3.6)$$

where $R = R(p_L, p)$ is determined from the $L_0 = 0$ condition (3.1). Note that the L_n and hence d are rotation/Lorentz invariant in $D-1$ dimension and that the result (3.6), as well as the d cohomology states, are, of course, independent of our special choice of coordinates.

It remains to consider the case $p = 0$. Then the above construction of $\hat{d}_0$ and K breaks down. We may now define $p^+(n) = \frac{1}{\sqrt{2}}(p_1 + i(p_L - (n+1)Q_L)) = \frac{i}{\sqrt{2}}(p_L - (n+1)Q_L)$ and $a_n^\pm = \frac{1}{\sqrt{2}}(a_n^1 \pm i a_n^L)$ and assign degrees again by eq. (3.3). Then

$$\hat{d}_0 = \sum_{n \neq 0} p^+(n) c_{-n} a_n^- . \quad (3.7)$$

If $p^+(n)$ is non-vanishing for all non-zero integers n , i.e. if $p_L \neq Q_L(r+1), r \in \mathbf{Z}(\neq 0)$, we

may construct the corresponding K operator ($K = \sum_{n \neq 0} \frac{1}{p^+(n)} b_n a_{-n}^+$) and conclude as above to obtain eq. (3.6).

However, for $p_L = Q_L(r + 1)$, we have to exclude the term $n = r$ from K and the $\hat{d}_0$ cohomology states now may contain b_r or a_r^- (for $r < 0$) and c_{-r} or a_{-r}^+ (for $r > 0$) [5] in addition to the $a_{-m}^i, i = 2, \dots, D$ oscillators. Thus, in general, the $\hat{d}_0$ cohomology is not confined to a single degree and the above theorem cannot be applied; we have to obtain the $\hat{d}$ cohomology in a different way. This will be the only case where we will go beyond the pure Fock space approach and use the vanishing theorem of ref. 4.

Note first that for $p = 0$, $p_L = Q_L(r + 1)$ the $L_0 = 0$ condition becomes

$$\frac{25 - D}{24}(r^2 - 1) = R - 1 . \quad (3.8)$$

Thus, for given D and r , there is not always an integer solution for the level R where non-trivial cohomology may occur, If there isn't we just have no non-trivial cohomology at all for the corresponding p_L . We see also from eq. (3.8) that (for $D \neq 1$) we always have $r^2 > R$, except for $r^2 = R = 1$. In the latter case $p_L = 0$ or $2Q_L$ and we know that the Liouville and matter Verma modules have null vectors $L_{-1}^L |p >$ and $L_{-1}^M |p >$ at level $r^2 = 1$ that vanish in the Fock space (or in the dual Fock space). So the level of the (vanishing) null vectors coincides with the level selected by the on-shell condition (3.8) and the vanishing theorem of ref. 4 does no longer apply (V is not a free $\mathcal{L}_-$ -module). New cohomology states with non-zero ghost-numbers are possible. Explicitly we find the cohomology states

$$\begin{aligned} a_{-1}^i |p_L = Q_L(1 + r), p = 0 > \quad r = \pm 1 , \\ b_{-1} |p_L = 0, p = 0 > \quad (r = -1) , \\ c_{-1} |p_L = 2Q_L, p = 0 > \quad (r = +1) . \end{aligned} \quad (3.9)$$

We see that with respect to the case of non-zero matter momentum or $p_L \neq Q_L(1 + r)$ we find an extra ghost-number zero and a ghost-number $r(= \pm 1)$ cohomology state.

For $r^2 > 1$ (and $1 < D < 25$),^{*} the level R selected by the $L_0 = 0$ condition (3.8) is always smaller than the level r^2 at which the Liouville null vector occurs. Thus, up to level $R < r^2$

^{*} Note that there is no possible integer solution R for $r = 0$ and the range of D we are considering.

there are no Liouville null vectors. This is enough to conclude that the combined Liouville plus matter system has no null vectors up to level R except when dealing with a state without Liouville excitations. Then we must still check for matter null states. In the matter sector, we have a representation of the Virasoro algebra with $1 < c_M < 25$ and highest weight $h = 0$. We know from the Kac determinant formula that, in this range of c , $h_{r's'}$ (with $r's' > 0$ since otherwise there are no null states) equals zero only for $r' = s' = \pm 1$, and the (matter) null vector is just the obvious $L_{-1}^M |p = 0\rangle$. The latter case corresponds to the extra cohomology states at level one discussed above (since these states are on shell only for $p_L = 0, 2Q_L$). Thus there are no null vectors of the combined Liouville plus matter system for $p = 0$ and $p_L = Q_L(r + 1)$ with $r \neq \pm 1$ at any level less than r^2 .

The BRST operator d does not change the (total) level R of a state. In particular, if we have a state ψ of level R , then the determination of whether $d\psi = 0$ or whether $\psi = d\chi$ only involves elements of the (matter plus Liouville) Verma module of levels R' with $R' \leq R$. Now, since there are no null vectors in the (matter plus Liouville) Verma module at any level $R' \leq R < r^2$, we conclude that we may apply the vanishing theorem of ref. 4.[†] As a result, for $r \neq \pm 1$ there is no non-trivial cohomology at non-zero ghost-number, and the dimensions of the zero ghost-number cohomology groups are again given by equation (3.6) where $R \equiv R(r)$ is given by the on-shell condition (3.8).

Finally, one might wish to consider the more general situation of D free matter fields one of which (called ϕ_M) has a background charge Q_M (real or imaginary). By a rotation in field space we may always reduce this to the case studied above with $\tilde{Q}_L = i \left(\frac{25-D}{12} \right)^{1/2}$ as before. Thus the cohomology is again described by eq. (3.6) except for the appearance of extra cohomology states (like (3.9)) at level one when $\tilde{p} = 0$ and $\tilde{p}_L = \tilde{Q}_L(1 + r)$ with $r = \pm 1$, which translates into $p_L = Q_L(1 + r)$ and $p_M = Q_M(1 + r)$ ($r = \pm 1$), and all other $p_i = 0$, in terms of the original (unrotated) momenta.

Acknowledgements:

It is a pleasure to acknowledge stimulating discussions with P. Bouwknegt and E. Witten.

[†] Originally this vanishing theorem has been proven for a free $\mathcal{L}_-$ -module ($\mathcal{L}_-$ being the negatively moded part of the Virasoro algebra). Our preceding argument just shows that it is sufficient for the cohomology at level R if there are no null vectors up to level R .

REFERENCES

1. M. Kato and K. Ogawa, Nucl. Phys. **B212** (1983) 443.
2. M.D. Freeman and D.I. Olive, Phys. Lett. **B175** (1986) 151.
3. M. Henneaux, Phys. Lett. **B177** (1986) 35.
4. I.B. Frenkel, H. Garland and G.J. Zuckerman, Proc. Natl. Acad. Sci. USA **83** (1986) 8442.
5. P. Bouwknegt, J. McCarthy and K. Pilch, CERN preprint CERN-TH.6162/91 (July 1991).
6. B.H. Lian and G.J. Zuckerman, Phys. Lett. **B254** (1991) 417, **B266** (1991) 21.
7. E. Witten, IAS preprint IASSNS-HEP-91/51 (August 1991).
8. D. Kutasov, E. Martinec and N. Seiberg, Princeton/Rutgers preprint PUPT-1293/RU-91-49 (November 1991);
I.R. Klebanov, Princeton preprint PUPT-1302 (December 1991).
9. E. Witten and B. Zwiebach, IAS preprint IASSNS-HEP-92/4 (January 1992).
10. J.-L. Gervais and A. Neveu, Phys. Lett. **B151** (1985) 271;
A. Bilal and J.-L. Gervais, Nucl. Phys. **B284** (1987) 397;
J.-L. Gervais, Phys. Lett. **B243** (1990) 85, Comm. Math. Phys. **138** (1991) 301.
11. P. Bouwknegt, J. McCarthy and K. Pilch, CERN preprint CERN-TH.6346/91 (November 1991).